

ON LEFT FBN RINGS¹

John A. Beachy²

Let R be a left Noetherian ring with identity. (All modules considered are unital left modules.) The ring R is said to be left FBN if for each prime ideal P of R , each left ideal of R/P that is essential in R/P contains a nonzero two-sided ideal. It is well known ([6], Proposition VII 2.4) that R is left FBN if for each finitely generated R -module M there exist $m_1, \dots, m_n \in M$ such that $\text{Ann}(M) = \text{Ann}(m_1, \dots, m_n)$, and Cauchon has shown in [3] that the converse is true. In this note we give a local version of the above result, and we show that R is left FBN if and only if this local condition holds for each minimal prime torsion theory.

For each R -module M , a torsion theory τ_M is defined in the following way: $\tau_M(X) = \{x \in X \mid f(x) = 0 \text{ for each } f \in \text{Hom}_R(X, E(M))\}$, where $E(M)$ denotes the injective envelope of M . Let σ be a torsion theory of $R\text{-Mod}$. Then an R -module X is called σ -torsion (σ -torsionfree) if $\sigma(X) = X$ ($\sigma(X) = 0$), and a submodule Y of X is called σ -dense (σ -closed) if X/Y is σ -torsion (σ -torsionfree). A torsion theory π is said to be prime if there exists a uniform R -module U such that $\pi = \tau_U$. In this case, if $\text{Ann}(x)$ is maximal in the set $\{\text{Ann}(u) \mid u \in U\}$, then each nonzero submodule of Rx is π -dense, and as a consequence the localization $(Rx)_\pi$ is a minimal subobject of U_π . If the ideal P is maximal in the set of annihilators of submodules of U , then P is a prime ideal of R , and $\tau_{R/P}$ is a prime in $R\text{-Mod}$ satisfying $\tau_{R/P} \geq \pi$ (there exists a uniform submodule A of R/P such that $\tau_{R/P} = \tau_A$ by Theorem 3.9 of [5]). The ideal P is called an associated prime ideal of U . We denote by $\text{ass}(M)$ the set of prime torsion theories π such that there exists a uniform submodule U of M such that $\pi = \tau_U$.

¹This is a translation of the original paper, which appeared in French.

²I would like to thank Professor J. Lambek and the Department of Mathematics of McGill University for their hospitality.

Lemma 1 *Let π be a prime torsion theory of $R\text{-Mod}$, and let M be a finitely generated R -module such that $\text{ass}(M) = \{\pi\}$. If $\{M_\alpha\}_{\alpha \in \Lambda}$ is a family of π -closed submodules of M such that $\bigcap_{\alpha \in \Lambda} M_\alpha = 0$, then there exists a finite subset Φ of Λ such that $\bigcap_{\alpha \in \Phi} M_\alpha = 0$.*

Proof. Since M is a Noetherian module and $\text{ass}(M) = \{\pi\}$, M contains an essential submodule $\bigoplus_{i=1}^n U_i$, such that each module U_i is uniform and defines π . Therefore $\pi M = 0$ and M_π contains an essential subobject $\bigoplus_{i=1}^n M_i$, such that each subobject M_i is minimal in M_π . The one-to-one correspondence ([6], Corollary IX 4.4) between the set of π -closed submodules of M and the set of subobjects of M_π (which preserves intersections) shows that $\bigcap_{\alpha \in \Lambda} (M_\alpha)_\pi = 0$, and so there exists a finite subset Φ of Λ such that $\bigcap_{\alpha \in \Phi} (M_\alpha)_\pi = 0$, and thus $\bigcap_{\alpha \in \Phi} M_\alpha = 0$. \square

Theorem 2 *If σ is a torsion theory of $R\text{-Mod}$, then the following conditions are equivalent.*

(1) *For each σ -closed prime ideal P of R , each essential σ -closed left ideal of R/P contains a nonzero two-sided ideal.*

(2) *For each finitely generated σ -torsionfree R -module M , there exist elements $m_1, \dots, m_n \in M$ such that $\text{Ann}(M) = \text{Ann}(m_1, \dots, m_n)$.*

Proof. (1) \Rightarrow (2). Let M be a finitely generated σ -torsionfree R -module. It is sufficient to establish condition (2) when M is a uniform R -module, since each finitely generated R -module has a decomposition $\bigcap_{i=1}^n X_i = 0$ such that X/X_i is uniform. Let $\pi = \pi_M$, $A = \text{Ann}(M)$, and $\rho \in \text{ass}(R/A)$. Then there exists a prime ideal P associated to ρ , and it is easy to verify that there exists a submodule N of M such that $P = \text{Ann}(N)$, since P is σ -closed in R . If we show that $\tau_{R/P} = \pi$, then $\rho = \pi$ and $\text{ass}(R/A) = \{\pi\}$, and it follows, from Lemma 1, that there exists a finite subset of the set $\{\text{Ann}(m) \mid m \in M\}$ such that $A = \bigcap_{i=1}^n \text{Ann}(m_i)$.

If $\text{Hom}_R(N', R/P) = 0$ for each submodule N' of N , then N is a singular R/P -module, and if $N = Rx_1 + \dots + Rx_k$, then $\bigcap_{i=1}^k \text{Ann}(x_i)/P$ is an essential σ -closed left ideal of R/P ; therefore it contains a nonzero two-sided ideal I/P , and $IN = 0$, which is impossible since $P = \text{Ann}(N)$. We can therefore conclude that there exists a submodule N' of N such that $\text{Hom}_R(N', R/P) \neq 0$, and if $f(N') \neq 0$, then there exist $y_1, \dots, y_m \in N'$ such that $P = \text{Ann}(f(y_1), \dots, f(y_m)) = \text{Ann}(y_1, \dots, y_m)$, since P is a prime ideal and there does not exist an infinite descending chain of annihilators of R/P . This shows that $E(R/P)$ is contained in a finite direct sum of copies

of $E(M)$, and Azumaya's theorem ([6], Proposition V 5.4) implies that there exists a submodule A of R/P such that $E(M) \simeq E(A)$. Therefore $\tau_{R/P} = \rho$.

(2) \Rightarrow (1). If P is a prime ideal of R and C/P is an essential σ -closed left ideal of R/P , then $R/\text{Ann}(R/C)$ is contained in a finite direct sum of copies of R/C . This shows that $P \subset \text{Ann}(R/C) \subseteq C$, since R/C is a singular R/P module but R/P is a nonsingular R/P -module. \square

If R satisfies the conditions of Theorem 2, then it is easy to check that the correspondence which associates to each indecomposable injective R -module E the unique prime ideal maximal in the set of annihilators of submodules of E induces a one-to-one correspondence between the set of isomorphism classes of σ -torsionfree indecomposable injective R -modules and the set of σ -closed prime ideals of R .

Corollary 3 *If σ is a maximal torsion theory of $R\text{-Mod}$, then for each finitely generated σ -torsionfree R -module there exist elements $m_1, \dots, m_n \in M$ such that $\text{Ann}(M) = \text{Ann}(m_1, \dots, m_n)$.*

Proof. If σ is maximal, then there exists a minimal prime ideal P such that $\sigma = \tau_{R/P}$ ([2], Theorem 4.6) and P is the only σ -closed prime ideal in R ([1], Proposition 1.2). If A/P is an essential left ideal of R/P , then $\text{Hom}_R(B/A, R/P) = 0$ for each left ideal B such that $A \subseteq B \subseteq R$, since B/A is a singular R/P -module, but R/P is a nonsingular R/P -module. Therefore A/P is not σ -closed in R/P , and thus condition (1) of Theorem 2 is trivially satisfied. \square

Theorem 4 *The ring R is left FBN if and only if each torsion theory minimal in the set of prime torsion theories of $R\text{-Mod}$ satisfies the conditions of Theorem 2.*

Proof. If R is left FBN, then the conditions of Theorem 2 are satisfied for every torsion theory.

Conversely, it is sufficient to show that for each finitely generated uniform R -module M there exist $m_1, \dots, m_n \in M$ with $\text{Ann}(M) = \text{Ann}(m_1, \dots, m_n)$. Let C be a left ideal of R maximal in the set of τ_M -closed left ideals of R . If π is prime in $R\text{-Mod}$ and π is strictly contained in τ_M , then C is strictly contained in a left ideal B maximal in the set of π -closed left ideals of R . Since R is left Noetherian, this construction eventually yields a torsion theory σ minimal in the set of prime torsion theories, and $\sigma \leq \tau_M$. Therefore $\sigma M \subseteq \tau_M(M) = 0$

and condition (2) of Theorem 2 show that there exist $m_1, \dots, m_n \in M$ such that $\text{Ann}(M) = \text{Ann}(m_1, \dots, m_n)$. \square

We note that if each prime torsion theory of $R\text{-Mod}$ is maximal, then it follows from Theorem 4 and Corollary 3 that R is left FBN. Therefore R is left Artinian, since each prime ideal of R is maximal ([6], Proposition VIII 1.14). This gives a new proof of Theorem 5.10 of Goldman [4].

References

- [1] BEACHY, J.A., On maximal torsion radicals, *Canad. J. Math.* **25**, 712–726 (1973).
- [2] BEACHY, J.A., On maximal torsion radicals II, *Canad. J. Math.* **27**, 115–120 (1975).
- [3] CAUCHON, G., Les T-anneaux et la condition de Gabriel, *C.R. Acad. Sci. Paris* **277**, 1153–1156 (1973).
- [4] GOLDMAN, O., Elements of noncommutative arithmetic I, *J. Algebra* **35**, 308–341 (1975).
- [5] LAMBEK, J. and MICHLER, G., The torsion theory at a prime ideal of a right Noetherian ring, *J. Algebra* **25**, 364–389 (1973).
- [6] B. STENSTRÖM, B. *Rings of Quotients* Berlin–Heidelberg–New York: Springer–Verlag, (1975).

McGill University
Montreal, Quebec
and
Northern Illinois University
DeKalb, Illinois